



Beefmaker Grower Pellet R20

(For beef cattle fed in confinement for slaughter)

Medicated

Improved feed efficiency.
ACTIVE DRUG INGREDIENT

Monensin (as Monensin Sodium) . 20 g/ton

Guaranteed Analysis

Crude Protein	Min	13.0 %	Potassium	Min	0.6 %
Crude Fat	Min	2.0 %	Zinc	Min	50 ppm
Crude Fiber	Max	16.5 %	Copper	Min	20 ppm
Calcium	Min	0.5 %	Iodine	Min	3.6 ppm
Calcium	Max	0.9 %	Selenium	Min	0.3 ppm
Phosphorus	Min	0.3 %	Vitamin A	Min	3,000 IU/lb
Salt	Min	0.5 %	Vitamin D	Min	300 IU/lb
Salt	Max	0.9 %	Vitamin E	Min	7 IU/lb

Ingredients

Cottonseed Hulls, Cotton Plant By-Product, Milo, Wheat Midds, Cottonseed Meal, Corn Gluten Feed, Corn Dried Distillers Grains Plus Solubles, Cane Molasses, Yeast Culture Concentrate, Saccharomyces Cerevisiae Fermentation Product, Magnesium Limestone, Calcium Carbonate, Salt, Lignin Sulfonate, Zinc Proteinates, Selenium Yeast, Manganese Sulfate, Zinc Sulfate, Zinc Oxide, Copper Chloride, Ferrous Sulfate, Ethylene Diamine Dihydrdide, Sodium Selenite, Calcium Iodate, Cobalt Carbonate, Vitamin A Acetate, Cholecalciferol (source of vitamin D3), Alpha Tocopherol Acetate (source of Vitamin E).

Feeding Directions

Feed to beef cattle at a rate of 15 to 25 lb to provide 150 to 250 mg Monensin per head per day. An unlimited supply of clean, fresh water should be available at all times. Contact your A-C representative for feeding and management recommendations specific to your operation.



NOTICE: Type C cattle feed containing 30g or less Monensin sodium per ton shall bear an expiration date of 30 days after the date of manufacture.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin could be fatal to cattle. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION: This product contains added copper. Do Not Use in feeds for Sheep.

CAUTION: Follow Label Directions. The addition of higher levels of selenium to feeds containing this product is not permitted.



Manufactured/Guaranteed By:
AC Nutrition, LP