



Ranch Forage RepX R20

(For goats)

Medicated

For the prevention of coccidiosis caused by Eimeria crandalis, Eimeria christenseni and Eimeria ninakohlyakimovae.

ACTIVE DRUG INGREDIENT

Monensin (as monensin sodium) . 20.0 g/ton

Guaranteed Analysis

Crude Protein	Min	10.0 %	Salt	Max	10.8 %
Crude Fat	Min	1.5 %	Potassium	Min	0.6 %
Crude Fiber	Max	20.0 %	Iodine	Min	2.4 ppm
Calcium	Min	0.5 %	Selenium	Min	0.2 ppm
Calcium	Max	0.9 %	Vitamin A	Min	2,500 IU/lb
Phosphorus	Min	0.3 %	Vitamin D	Min	250 IU/lb
Salt	Min	9.0 %	Vitamin E	Min	5 IU/lb

Ingredients

Corn Gluten Feed, Cottonseed Hulls, Cotton Plant By-Product, Rice Hulls, Milo, Wheat Midds, Corn Dried Distillers Grains Plus Solubles, Pinto Beans, Cane Molasses, Calcium Carbonate, Salt, Monocalcium Phosphate, Zinc Proteinates, Selenium Yeast, Manganese Sulfate, Zinc Sulfate, Zinc Oxide, Ferrous Sulfate, Ethylene Diamine Dihydriodide. Sodium Selenite, Calcium Iodate, Cobalt Carbonate, Vitamin A Acetate, Cholecalciferol (source of vitamin D3), Alpha Tocopherol Acetate (source of Vitamin E).

Feeding Directions

Feed to goats continuously as their sole diet. An unlimited supply of clean, fresh water should be available at all times. Contact your A-C representative for feeding and management recommendations specific to your operation. An unlimited supply of clean, fresh water should be available at all times. Contact your A-C representative for feeding and management recommendations specific to your operation.



NOTICE: Type C cattle feed containing 30g or less Monensin sodium per ton shall bear an expiration date of 30 days after the date of manufacture.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin could be fatal to cattle. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION: Follow Label Directions. The addition of higher levels of selenium to feeds containing this product is not permitted.



Manufactured/Guaranteed By:
AC Nutrition, LP

Expiration Date:

Expira